

Manitoba Weekly

West Nile virus

Surveillance Report

Week 33 – (August 12 - 18, 2018)

Communicable Disease Control

Public Health Branch

Active Living, Indigenous Relations, Population &

Public Health Care Division

Manitoba Health, Seniors and Active Living

Released: August 24, 2018

About the Surveillance Report

The weekly 'West Nile Virus Surveillance Report' outlines the most current surveillance data and is posted weekly on the website (www.gov.mb.ca/health/wnv) during the summer season. Surveillance data are subject to change and will be updated accordingly as new information becomes available.

Manitoba Health, Seniors and Active Living (MHSAL) conducts surveillance for West Nile virus (WNV) within human, mosquito & horse populations annually:

- **Mosquito:** Mosquito surveillance is conducted twice per week between mid-May and mid-September (weather dependent) in a number of southern Manitoba communities. In Manitoba WNV testing is conducted on *Culex tarsalis* mosquitoes, the principal vectors of WNV, and both mosquito numbers and infection rates (i.e. positive mosquito pools*) are reported.
 - Communities chosen for mosquito trap placement were selected based on population density, local evidence of prior WNV activity and representative geographic distribution.
- **Human:** Human WNV surveillance is conducted throughout the year (January – December) by Cadham Provincial Laboratory and Canadian Blood Services, with all data reportable to MHSAL.
 - Human cases are included in the Weekly WNV Surveillance Report based on the date they are reported to MHSAL. Case classification information is not included in this report but can be found on the website (www.gov.mb.ca/health/wnv/stats.html).
- **Horse:** Surveillance of WNV in horses is conducted by Manitoba Agriculture with cases reported to MHSAL as detected.

The risk of WNV transmission is expected to be present throughout southern Manitoba each year and mosquito trapping provides a localized estimate of WNV risk. The absence of traps in a community or region does not imply that there is no risk of WNV in those locations. Further, low *Culex tarsalis* numbers and/ or infection rates should not be interpreted as zero risk. Residents and visitors are strongly encouraged to protect themselves from mosquito bites throughout the season even in areas with no mosquito traps or low WNV activity.

The accumulation of Degree Days¹ are recorded throughout the season as there is a general correlation between increased and/ or rapid accumulation of Degree Days and WNV transmission risk. Warmer temperatures associated with increased Degree Days serve to decrease mosquito development times, shorten the WNV incubation period and increase biting activity. All of which can increase the risk of WNV transmission, should other conditions also be favourable. Seasonally the greatest accumulation of Degree Days typically occurs in the southwestern portion of the province and along the Red River valley.

For additional West Nile virus information, including precautionary measures and symptoms, please consult the MHSAL WNV website (www.gov.mb.ca/health/wnv) or contact Health Links at 204-788-8200 (in Winnipeg) or toll free at 1-888-315-9257.

¹ For more detailed description of mosquito pools and degree days please consult **Appendix 2**.

WNV Provincial Surveillance Data

- During Week 33* (August 12 - 18) MHSAL detected six (6) additional human WNV cases, bringing the season total to date to seven (7).
 - One case has been classified as WNV Neurological Syndrome, one as WNV Non-Neurological Syndrome, three as Asymptomatic, while the classification of two cases remain under investigation pending additional information.
- During Week 33, MHSAL also detected an additional twenty-three (23) WNV positive *Culex tarsalis* mosquito pools, bringing the total to date this season to one-hundred and fifty-two (152) (Figure 1 & 3, Table 1).
 - The positive pools were collected from Interlake-Eastern (Beausejour and Stonewall), Prairie Mountain (Boissevain, Carberry, Killarney, Sioux Valley and Virden), Southern (Portage la Prairie and Winkler) and Winnipeg (West St. Paul and Winnipeg) Health Regions.
- In Week 33 *Cx. tarsalis* activity was recorded in all four southern Manitoba Health Regions. Activity increased slightly in comparison to Week 32 and specimens were collected from all of the 29 sentinel communities (Figure 1, Table 1).
 - *Cx. tarsalis* activity was greatest in the Prairie Mountain Health Region in Week 33 (108.19 *Cx. tarsalis*/ trap night). Infection rates were also highest in the Prairie Mountain Health Region (17.8%).
- In addition, fourteen (14) WNV positive birds (Interlake-Eastern and Winnipeg Health Regions) and four WNV positive horses (Interlake-Eastern Health Region and Prairie Mountain Health) have also been reported this season.

* For a listing of CDC surveillance weeks and corresponding dates for 2018 please see Appendix 1.

Table 1 – Average number of *Culex tarsalis* mosquitoes captured by Health Region (current to Week 33)

Health Region	CDC Week								
	25	26	27	28	29	30	31	32	33
Interlake-Eastern	8.16	7.00	23.47	54.88	182.65	58.84	19.58	23.74	20.22
Prairie Mountain	1.12	7.74	11.26	65.50	139.15	80.83	20.60	57.57	108.19
Southern	9.34	23.41	20.75	133.44	57.70	84.64	23.60	31.20	20.48
Winnipeg	17.76	5.91	12.06	75.47	64.11	96.94	34.98	26.50	25.11
Provincial Average	8.42	12.36	16.19	89.82	101.60	83.17	26.10	36.96	48.35
Historical Avg	10.60	108.79	149.49	132.39	99.27	234.27	251.73	118.00	68.70
	Indicates that one or more positive mosquito pools were detected within the health region.								




Figure 1 – WNV activity by Health Region within Manitoba (current to Week 33).

Table 2 – Average number and proportion of *Culex tarsalis* mosquitoes collected by surveillance community* in southern Manitoba – three week trend (current to Week 33).

Health Region	Community	Week 33		Week 32	
		Avg # of <i>Cx. tarsalis</i>	Proportion of <i>Cx. tarsalis</i>	Avg # of <i>Cx. tarsalis</i>	Proportion of <i>Cx. tarsalis</i>
Interlake-Eastern	Beausejour	38.00	83.52	9.00	55.10
	Gimli	19.25	27.11	22.50	60.00
	Oakbank	2.00	1.83	4.50	10.34
	Selkirk	9.75	25.00	54.25	57.71
	Stonewall	30.00	63.38	24.75	14.39
Prairie Mountain	Boissevain	687.50	96.83	250.25	86.14
	Brandon	15.50	68.89	17.80	58.94
	Carberry	30.75	73.21	24.25	72.39
	Dauphin	3.25	11.30	1.00	3.92
	Killarney	134.00	94.37	58.75	55.56
	Minnedosa	5.50	4.92	2.50	1.43
	Sioux Valley FN	138.25	60.44	116.00	31.56
	Souris	31.75	44.41	21.50	67.19
Southern	Virden	66.25	86.60	85.75	88.86
	Altona	17.00	37.99	59.75	51.73
	Carman	6.50	27.37	7.25	22.31
	Headingley	1.00	100.00	1.00	16.67
	Morden	29.75	58.91	73.75	34.58
	Morris	4.50	25.00	13.75	4.93
	Niverville	32.67	97.03	25.75	23.79
	Portage la Prairie	62.50	73.96	25.25	38.55
	Roseau River FN	1.75	6.73	40.75	45.92
	Ste. Anne	5.75	28.05	2.75	13.41
	Sandy Bay FN	12.75	24.52	1.25	1.97
	Steinbach	13.33	100.00	3.50	48.28
Winkler	53.00	30.75	97.00	33.42	
Winnipeg	East St Paul	7.00	33.33	3.00	3.87
	West St Paul	78.50	84.41	128.50	39.66
	Winnipeg	22.91	54.02	21.27	23.69
	Indicates that one or more positive mosquito pools were detected within the community.				

* Top three communities with the highest weekly average of *Culex tarsalis* are indicated in bold.




Figure 2 – Average number of *Culex tarsalis* mosquitoes collected across southern Manitoba during Week 33.





Figure 3 – Distribution of WNV positive *Culex tarsalis* mosquito pools collected in southern Manitoba (current to Week 33).



Source: Map produced courtesy of Agriculture and Agri-Food Canada's Prairie Pest Monitoring Network.

Figure 4 - Degree day accumulations, as of Week 33, across the Prairie Provinces

Table 3 – Total number of human WNV cases*, by Health Region of residence, reported to Manitoba Health, Seniors and Active Living by laboratories (current to Week 33).

Health Region	CDC Week									Totals
	25	26	27	28	29	30	31	32	33	
Interlake-Eastern	0	0	0	0	0	0	0	0	3	3
Prairie Mountain	0	0	0	0	0	0	0	0	2	2
Southern	0	0	0	0	0	1	0	0	0	1
Winnipeg	0	0	0	0	0	0	0	0	1	1
Totals	0	0	0	0	0	1	0	0	6	7

* Note that cases are presented by week reported to MHSAL, adjustments may be made as more details (such as exposure CDC week) become available through follow-up investigation.

Table 4 – Total number of *Culex tarsalis* mosquito pools tested during the 2018 season by health region (current to Week 33)

Health Region	CDC Week									Totals
	25	26	27	28	29	30	31	32	33	
Interlake-Eastern	19	16	20	23	41	30	19	20	14	214
Prairie Mountain	20	31	36	55	78	62	43	56	73	495
Southern	40	53	42	106	73	88	45	57	39	587
Winnipeg	28	24	24	57	48	67	62	39	37	432
Weekly Totals	107	124	122	241	240	247	169	172	163	1728

Table 5* – Total number and percentage of WNV positive *Culex tarsalis* mosquito pools by Health Region (current to Week 33)

Health Region	CDC Week									Totals
	25	26	27	28	29	30	31	32	33	
Interlake-Eastern	0 (0)	0 (0)	3 (15.0)	0 (0)	6 (14.6)	7 (23.3)	2 (10.5)	0 (0)	2 (14.3)	20 (9.3)
Prairie Mountain	0 (0)	0 (0)	0 (0)	2 (3.6)	5 (6.4)	10 (16.1)	6 (14.0)	8 (14.3)	13 (17.8)	44 (8.9)
Southern	0 (0)	0 (0)	0 (0)	5 (4.7)	8 (11.0)	13 (14.8)	4 (8.9)	13 (22.8)	2 (5.1)	45 (7.7)
Winnipeg	1 (3.6)	0 (0)	0 (0)	4 (7.0)	6 (12.5)	11 (16.4)	9 (14.5)	6 (15.4)	6 (16.2)	43 (10.0)
Weekly Totals	1 (0.9)	0 (0)	3 (2.5)	11 (4.6)	25 (10.4)	41 (16.6)	21 (12.4)	27 (15.7)	23 (14.1)	152 (8.8)

* Note that numbers outside brackets represent positive pools, numbers within represent the percentage of total pools that tested positive for WNV.

Table 6 – Comparison of year-to-date cumulative and year-end total West Nile virus in Manitoba (current to Week 33)

Year	Cumulative (Year-to-Date) Amount		Year End Totals	
	Positive Mosquito Pools	Human WNV Cases	Positive Mosquito Pools	Human WNV Cases
2018	152	7	TBD	TBD
2017	37	4	41	5
2016	37	18	39	24
2015	28	4	30	5
2014	14	4	24	5
2013	17	3	19	3
2012	108	33	116	39
2011	0	0	0	0
2010	19	0	20	0
2009	2	2	2	2
2008	31	10	41	12
2007	894	431	948	587
2006	161	41	171	51
2005	187	40	193	58
2004	54	3	57	3
2003	149	66	290	143

WNV Activity in Canada and the United States

Canada:

- As of Week 33 there have been 16 WNV human cases (7 in Manitoba and 9 in Ontario including 2 travel-related cases), 309 WNV positive mosquito pools (153 in Manitoba, 115 in Ontario, 8 in Quebec and 33 in Saskatchewan), 59 WNV positive birds (14 in Manitoba, 15 in Ontario, 7 in Quebec and 2 in Saskatchewan) and 11 WNV positive horses (4 in Manitoba, 2 in Saskatchewan and 5 in Alberta) reported in Canada. There have been no other positive WNV surveillance indicators reported in Canada to date.
- Additional Canadian WNV information can be obtained by consulting the Public Health Agency of Canada West Nile virus website at <https://www.canada.ca/en/public-health/services/diseases/west-nile-virus/surveillance-west-nile-virus.html>, or by consulting the respective provincial department websites.

United States:

- As of August 21, 2018 the US CDC is reporting WNV activity from 45 states (includes human cases, positive mosquito pools and positive birds). A total of 231 human cases along with 89 presumptive blood donors have been reported to date, of which 58% have been classified as West Nile neuroinvasive disease. The US CDC is also reporting eight WNV related deaths this year.
 - WNV activity has been reported from Minnesota (20 human cases (including 16 presumptive viremic blood donors) and numerous positive mosquito pools), North Dakota (18 human cases, 34 WNV positive mosquito pools, 1 WNV positive horse and 10 WNV positive birds) and South Dakota (52 human cases (including 2 WNV associated mortality and 12 presumptive viremic blood donors) and numerous mosquito pools).
- Up to date U.S. WNV information can be obtained by visiting the United States Centers for Disease Control and Prevention – West Nile virus Website' at <https://www.cdc.gov/westnile/statsmaps/preliminarymapsdata2018/index.html>, or by consulting state specific Public Health websites.

Appendix 1

Table 8 – 2018 CDC surveillance weeks

CDC Week Number	Dates	CDC Week Number	Dates
21	May 20 – May 26	30	July 22 - July 28
22	May 27 – June 2	31	July 29 - August 4
23	June 3 - June 9	32	August 5 - August 11
24	June 10 - June 16	33	August 12 - August 18
25	June 17 - June 23	34	August 19 - August 25
26	June 24 – June 30	35	August 26 - September 1
27	July 1 - July 7	36	September 2 - September 8
28	July 8 - July 14	37	September 9 - September 15
29	July 15 - July 21	38	September 16 - September 22

Appendix 2

Average number of *Culex tarsalis* – This weekly value provides an estimate of the *Culex tarsalis* numbers and activity. The potential risk of WNV transmission is greater when more *Culex tarsalis* are present – should the virus itself be present and other conditions prove favorable. It is calculated by dividing the total number of *Culex tarsalis* mosquitoes captured in the specified area by the total number of trap nights for the week (a trap night is recorded for each night that a trap was operational).

EXAMPLE: 120 *Culex tarsalis* collected; 2 traps operating on 2 nights (= 4 trap nights);
Average number = 120 (*Culex tarsalis*) / 4 trap nights = 30.0

Degree Day – Degree days are a measurement of heat accumulation. The threshold temperature below which West Nile virus development does not occur (when in mosquitoes) is 14.3°C. Degree days are calculated by taking the daily mean temperature and subtracting the cut-off threshold:

EXAMPLE: Mean Temperature = 19.3°C; Degree Day threshold = 14.3°C; 19.3 – 14.3 = 5.0 Degree Days.

During the season a running total of accumulated Degree Days is recorded. It is generally assumed that a total of 109 Degree Days are required for virus development to be completed and potential transmission to occur. The risk of transmission increases with increasing Degree Day accumulation. Moreover, consistently warmer temperatures will significantly shorten virus development time thereby increasing the potential risk of WNV transmission – should the virus itself be present and other conditions prove to be favorable.

Mosquito Pool – Mosquitoes of the same species, collected from the same trap on the same date are pooled together for the purposes of laboratory testing. *Culex tarsalis* mosquitoes collected from one trap on a given night are placed in pools of 1 – 50 mosquitoes for WNV testing. When more than 50 *Culex tarsalis* mosquitoes are collected from the same trap multiple pools are tested. Thus a positive pool refers to the detection of WNV in between 1 – 50 *Culex tarsalis* mosquitoes collected from a given trap.